

```
=> file casreact
FILE 'CASREACT' ENTERED AT 16:04:42 ON 10 JUN 2008
USE IS SUBJECT TO THE TERMS OF YOUR CUSTOMER AGREEMENT
COPYRIGHT (C) 2008 AMERICAN CHEMICAL SOCIETY (ACS)
```

Copyright of the articles to which records in this database refer is held by the publishers listed in the PUBLISHER (PB) field (available for records published or updated in Chemical Abstracts after December 26, 1996), unless otherwise indicated in the original publications.

FILE CONTENT:1840 - 7 Jun 2008 VOL 148 ISS 24


New CAS Information Use Policies, enter HELP USAGETERMS for details.

```
*****
*
*   CASREACT now has more than 13.8 million reactions   *
*
*****
```

Some CASREACT records are derived from the ZIC/VINITI database (1974-1999) provided by InfoChem, INPI data prior to 1986, and Biotransformations database compiled under the direction of Professor Dr. Klaus Kieslich.

This file contains CAS Registry Numbers for easy and accurate substance identification.

```
=> d que
L1          STR
```




Structure attributes must be viewed using STN Express query preparation.
L3 3 SEA FILE=CASREACT SSS FUL L1 (5 REACTIONS)

```
=> d l3 1-3 ibib abs hit
```

```
L3  ANSWER 1 OF 3  CASREACT  COPYRIGHT 2008 ACS on STN
ACCESSION NUMBER:  141:395464  CASREACT
TITLE:             Synthesis and Conformational Analysis of a Non-Amidine
                   Factor Xa Inhibitor That Incorporates
                   5-Methyl-4,5,6,7-tetrahydrothiazolo[5,4-c]pyridine as
                   S4 Binding Element
AUTHOR(S):         Haginoya, Noriyasu; Kobayashi, Syozo; Komoriya,
                   Satoshi; Yoshino, Toshiharu; Suzuki, Makoto; Shimada,
                   Takashi; Watanabe, Kengo; Hirokawa, Yumiko; Furugori,
                   Taketoshi; Nagahara, Takayasu
CORPORATE SOURCE:  Medicinal Chemistry Research Laboratory, Daiichi
                   Pharmaceutical Co. Ltd, Edogawa-ku, Tokyo, 134-8630,
                   Japan
SOURCE:            Journal of Medicinal Chemistry (2004), 47(21),
                   5167-5182
```

PUBLISHER:
DOCUMENT TYPE:
LANGUAGE:
GI

CODEN: JMCMAR; ISSN: 0022-2623
American Chemical Society
Journal
English




I

AB Our exploratory study was based on the concept that a non-amidine factor Xa (fXa) inhibitor is suitable for an orally available anticoagulant. We synthesized and evaluated a series of N-(6-chloronaphthalen-2-yl)sulfonylpiperazine derivs. incorporating various fused-bicyclic rings containing an aliphatic amine expected to be S4 binding element. Among this series, 5-methyl-4,5,6,7-tetrahydrothiazolo[5,4-c]pyridine type I displayed orally potent anti-fXa activity and evident prolongation of prothrombin time (PT) with the moderate bioavailability in rats. The X-ray crystal anal. afforded an obvious binding mode that 5-methyl-4,5,6,7-tetrahydrothiazolo[5,4-c]pyridine and 6-chloronaphthalene resp. bound to S4 and S1 subsites. In this X-ray study, we discovered a novel intramol. S-O close contact. Ab initio energy calcns. of model compds. deduced that conformers with the most close S-O proximity were most stable. The Mulliken population anal. proposed that this energy profile was caused by both of electrostatic S-O affinity and N-O repulsion. The results of these calcns. and X-ray anal. suggested a possibility that the restricted conformation effected the affinity to S4 subsite of fXa.

REFERENCE COUNT: 72 THERE ARE 72 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT


RX(150) OF 352 COMPOSED OF RX(23), RX(24), RX(25)
RX(150) BY + BZ + B + BR ==> CD



BY



BZ




B



BR

3
STEPS
→

10/578,844



● Li

CD

YIELD 82%

RX(23) RCT BY 89424-04-4, BZ 115-08-2
PRO CA 165948-22-1
SOL 71-36-3 BuOH
CON 2.5 hours, 100 deg C
NTE molecular sieves used

RX(24) RCT CA 165948-22-1

STAGE(1)

RGT N 1310-73-2 NaOH
SOL 7732-18-5 Water
CON SUBSTAGE(1) 2 hours, 110 deg C
SUBSTAGE(2) 110 deg C -> room temperature

STAGE(2)

RCT B 24424-99-5
SOL 67-56-1 MeOH
CON 2 hours, room temperature

STAGE(3)

RGT AH 7647-01-0 HCl
SOL 7732-18-5 Water
CON room temperature, pH 2 - 3

PRO CC 165948-24-3

RX(25) RCT CC 165948-24-3

STAGE(1)

RGT BT 109-72-8 BuLi
SOL 60-29-7 Et2O, 110-54-3 Hexane
CON 15 minutes, -78 deg C

STAGE(2)

RCT BR 124-38-9
CON 5 minutes, -78 deg C

PRO CD 365996-70-9

TITLE: Facile methods for preparation of thiazolopyridine and tetrahydrothiazolopyridine derivatives

AUTHOR(S): Haginoya, Noriyasu; Komoriya, Satoshi; Osanai, Ken; Yoshino, Toshiharu; Nagata, Tsutomu; Nagamochi, Masatoshi; Muto, Ryo; Yamaguchi, Mitsuhiko; Nagahara, Takayasu; Kanno, Hideyuki

CORPORATE SOURCE: Medicinal Chemistry Research Laboratory, Daiichi Pharmaceutical Co., Ltd, Tokyo, 134-8630, Japan

SOURCE: Heterocycles (2004), 63(7), 1555-1561
CODEN: HTCYAM; ISSN: 0385-5414



PUBLISHER: Japan Institute of Heterocyclic Chemistry

DOCUMENT TYPE: Journal

LANGUAGE: English

AB Improved routes to prepare tetrahydrothiazolo[5,4-c]pyridine-2-carboxylic acid lithium salts were developed. Route A consisted of the improved preparation of thiazolopyridine intermediates, and Route B is applicable for a large scale synthesis of tetrahydrothiazolo[5,4-c]pyridine-2-carboxylic acid derivs. The methods may serve as facile means for preparing thiazolopyridine and tetrahydrothiazolopyridine derivs.

RX(31) OF 33 COMPOSED OF RX(9), RX(10), RX(13)
RX(31) V + W + S ==> AO



AO
YIELD 66%

RX(9) RCT V 79099-07-3

STAGE(1)
RGT Y 123-75-1 Pyrrolidine

10/578,844

CAT 104-15-4 TsOH
SOL 110-82-7 Cyclohexane
CON 2 hours, reflux

STAGE(2)

RCT W 420-04-2
RGT D 10544-50-0 S8
SOL 67-56-1 MeOH
CON SUBSTAGE(1) 0 deg C
SUBSTAGE(2) 5 hours, 0 deg C

PRO X 365996-05-0
NTE scalable, >100 g

RX(10)

RCT X 365996-05-0
RGT AD 540-80-7 t-BuONO, AE 7789-45-9 CuBr2
PRO AC 365996-06-1
SOL 68-12-2 DMF
CON SUBSTAGE(1) 50 deg C
SUBSTAGE(2) 2 hours, 50 - 60 deg C
NTE scalable, >100 g

RX(13)

RCT AC 365996-06-1

STAGE(1)


RGT C 109-72-8 BuLi
SOL 60-29-7 Et2O, 110-54-3 Hexane
CON SUBSTAGE(1) -78 deg C
SUBSTAGE(2) 20 minutes, -78 deg C

STAGE(2)

RCT S 124-38-9
CON SUBSTAGE(1) 5 minutes, -78 deg C
SUBSTAGE(2) -78 deg C -> room temperature

PRO AO 365996-70-9

RX(33) OF 33 COMPOSED OF RX(9), RX(10), RX(11), RX(12)
RX(33) V + W + AF + S ==> U



V



W




AF



S

4
STEPS
→

10/578,844



● Li

U

YIELD 99%

RX(9) RCT V 79099-07-3

STAGE(1)

RGT Y 123-75-1 Pyrrolidine

CAT 104-15-4 TsOH

SOL 110-82-7 Cyclohexane

CON 2 hours, reflux

STAGE(2)

RCT W 420-04-2

RGT D 10544-50-0 S8

SOL 67-56-1 MeOH

CON SUBSTAGE(1) 0 deg C

SUBSTAGE(2) 5 hours, 0 deg C

PRO X 365996-05-0

NTE scalable, >100 g

RX(10)

RCT X 365996-05-0

RGT AD 540-80-7 t-BuONO, AE 7789-45-9 CuBr₂

PRO AC 365996-06-1

SOL 68-12-2 DMF

CON SUBSTAGE(1) 50 deg C

SUBSTAGE(2) 2 hours, 50 - 60 deg C

NTE scalable, >100 g

RX(11)

RCT AC 365996-06-1

STAGE(1)

RGT AH 76-05-1 F3CCO₂H

SOL 75-09-2 CH₂Cl₂

CON 10 minutes, room temperature

STAGE(2)

RCT AF 50-00-0

RGT AI 56553-60-7 Na.(AcO)₃BH, AJ 121-44-8 Et₃N, AK 64-19-7 AcOH

SOL 7732-18-5 Water, 75-09-2 CH₂Cl₂

CON SUBSTAGE(1) room temperature

SUBSTAGE(2) 1 hour, room temperature

STAGE(3)

10/578,844

RGT AL 1310-73-2 NaOH
SOL 7732-18-5 Water
CON room temperature

PRO AG 143150-92-9
NTE scalable, 50 g

RX(12) RCT AG 143150-92-9

STAGE(1)

RGT C 109-72-8 BuLi
SOL 60-29-7 Et2O, 110-54-3 Hexane
CON SUBSTAGE(1) -78 deg C
SUBSTAGE(2) -78 deg C -> 0 deg C
SUBSTAGE(3) 20 minutes, 0 deg C
SUBSTAGE(4) 0 deg C -> -78 deg C


STAGE(2)

RCT S 124-38-9
CON SUBSTAGE(1) 5 minutes, -78 deg C
SUBSTAGE(2) -78 deg C -> room temperature

PRO U 259809-25-1
NTE scalable, 50 g

L3 ANSWER 3 OF 3 CASREACT COPYRIGHT 2008 ACS on STN

ACCESSION NUMBER: 141:123587 CASREACT
TITLE: Orally active factor Xa inhibitors:
4,5,6,7-tetrahydrothiazolo[5,4-c]pyridine derivatives
AUTHOR(S): Haginoya, Noriyasu; Kobayashi, Syozo; Komoriya,
Satoshi; Hirokawa, Yumiko; Furugori, Taketoshi;
Nagahara, Takayasu
CORPORATE SOURCE: Medicinal Chemistry Research Laboratory, Daiichi
Pharmaceutical Co. Ltd., Edogawa-ku, Tokyo, 134-8630,
Japan
SOURCE: Bioorganic & Medicinal Chemistry Letters (2004),
14(11), 2935-2939
CODEN: BMCLE8; ISSN: 0960-894X
PUBLISHER: Elsevier Science B.V.
DOCUMENT TYPE: Journal
LANGUAGE: English
GI




AB In an investigation of factor Xa inhibitors, a series of
1-(6-chloronaphthalen-2-yl)sulfonyl-4-(4,5,6,7-tetrahydrothiazolo[5,4-
c]pyridine-2-carbonyl)piperazines were synthesized. In vitro inhibitory
activities of the compds. against factor Xa and coagulation are

summarized. Among these, 4-[(6-chloro-2-naphthalenyl)sulfonyl]-1-[(4,5,6,7-tetrahydro-5-methylthiazolo[5,4-c]pyridin-2-yl)carbonyl]-2-piperazinecarboxamide (I) and 4-[(6-chloro-2-naphthalenyl)sulfonyl]-N-methyl-1-[(4,5,6,7-tetrahydro-5-methyloxazolo[5,4-c]pyridin-2-yl)carbonyl]-2-piperazinecarboxamide, possessing a carbamoyl or N-methylcarbamoyl moiety, showed potent inhibitory activities when administered orally to rats.

REFERENCE COUNT: 18 THERE ARE 18 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT


RX(61) OF 188 COMPOSED OF RX(3), RX(4), RX(1)
RX(61) L + M + P + B ==> C



L



M




P



B

3
STEPS
→



● Li

C
YIELD 85%

RX(3) RCT L 89424-04-4, M 115-08-2
PRO N 165948-22-1
SOL 64-17-5 EtOH
NTE 4Å MS used

RX(4) RCT N 165948-22-1

STAGE(1)
RGT Q 121-44-8 Et3N
SOL 7732-18-5 Water

10/578,844

STAGE(2)

RCT P 24424-99-5

PRO A 165948-24-3

RX(1) RCT A 165948-24-3

STAGE(1)

RGT D 109-72-8 BuLi

SOL 60-29-7 Et2O


STAGE(2)

RCT B 124-38-9

PRO C 365996-70-9

RX(62) OF 188 COMPOSED OF RX(3), RX(5), RX(25)

RX(62) L + M + B ==> F



L




M



B

3
STEPS
→



● Li

F
YIELD 100%

RX(3) RCT L 89424-04-4, M 115-08-2
PRO N 165948-22-1
SOL 64-17-5 EtOH
NTE 4Å MS used

RX(5) RCT N 165948-22-1
RGT T 16853-85-3 LiAlH4
PRO S 259809-24-0
SOL 60-29-7 Et2O

10/578,844

RX(25) RCT S 259809-24-0

STAGE(1)

RGT D 109-72-8 BuLi

SOL 60-29-7 Et2O

STAGE(2)

RCT B 124-38-9

PRO F 259809-25-1

=>